

# **Monthly report on livestock disease trends as informally reported by veterinarians belonging to the Ruminant Veterinary Association of South Africa (RuVASA), a group of the South African Veterinary Association**

## **August 2018**

(Previous disease reports can be seen on the RuVASA website [www.ruvasa.co.za](http://www.ruvasa.co.za) )

These reports include data from individual practices

Click on Disease Reports

**The following practices and laboratories (120) submitted reports during August 2018:**

### **Mpumalanga (9)**

Balfour – Dr. Louis van Jaarsveld

Bethal – Dr. Hardus Pieters

Ermelo – Dr. Ben Potgieter

Grootvlei – Dr. Neels van Wyk

Karino – Dr. Silke Pfitzer

Middelburg – Drs. Erasmus, Malan and Bernitz

Nelspruit – Dr. André Beytel

Piet Retief – Drs. Niebuhr and Weber

Volksrust – Dr. Johan Blaauw

### **Gauteng (7)**

Bronkhorstspuit – Dr. De Bruin, De Bruin and Labuschagne

Hammanskraal – Dr. Hentie Engelbrecht

Magaliesburg – Dr. Ryan Jeffery

Nigel – Dr. Cindy van der Westhuizen

Onderstepoort Veterinary Academic Hospital – Proff. Annandale, Shakespear, Holm, Pettey and Drs, Fitte, Grobler, Hamman, Koeppel, Leask, Mabui, Marufu, Mokoele, O'Dell, Tshuma and Van der Leek

Pretoria – Dr. Hanneke Pienaar

Vanderbijlpark – Dr. Kobus Kok

### **Limpopo (9)**

Bela-Bela – Dr. Nele Sabbe

Makhado (Louis Trichardt) – Drs. Harris, Kilian, Bester and Herbst

Modimolle (Nylstroom) – Drs. Huber, Bredell and Barnard

Mokopane (Potgietersrust)- Dr. Henk Visser

Mokopane - Dr. Alwyn Venter (CCS)

Polokwane (Pietersburg) – Drs. Watson, Viljoen, Jansen van Vuuren, Van Rooyen, Snyman and Cremona

Tzaneen – ZZ2 Farm practice – Dr. Danie Odendaal

Vaalwater – Dr. Hampie van Staden

Vaalwater – Dr. Annemieke Müller

### **North West (10)**

Brits – Dr. Boshoff and Coertze

Christiana - Dr. Pieter Nel

Klerksdorp – Drs. Geral, Theron, Van den Berg and Van den Berg

Klerksdorp – Drs. Coetzee and Venter

Leeudoringstad – Dr. Ian Jonker

Lichtenburg – Dr. Nelmarie -Krüger-Rall

Rustenburg – Drs. Grobler, Sparks, Van Egdom, Van Rooyen, Goosen and Van Rensburg

Stella - Dr. Magdaleen Vosser

Ventersdorp/ Koster –Drs. Benadé and Van der Merwe

Vryburg – Dr. Jurie Kritzing

### **Free State (20)**

Bethlehem – Drs. Strydom and Strydom

Clocolan – Drs. Wasserman and Basson

Dewetsdorp – Dr. Marike Badenhorst

Excelsior/Ladybrand – Drs. Dedré Nel and De Vos

Ficksburg – Drs. Kotzé and Coetzer

Frankfort - Drs. Lessing, Cilliers and Janse van Rensburg

Gariep Dam – Dr. Marni Malan Strauss

Hertzogville – Dr. Nico Hendrikz

Hoopstad – Dr. Kobus Pretorius

Kroonstad – Drs. Daffue, Eksteen, Van Zyl and Van der Walt

Memel – Drs. Nixon and Nixon

Parys – Drs. Wessels and Wessels

Philippolis – Dr. Stephan van Niekerk

Reitz - Dr. Murray Smith

Smithfield – Dr. Nienke van Hasselt

Viljoenskroon – Dr. Johan Kahts

Villiers – Drs. Hattingh and Hauptfleisch

Vrede – Drs. Myburgh, Bester-Cloete and Bester

Wesselsbron – Dr. Johan Jacobs

Zastron – Drs. Troskie and Strauss

**KwaZulu-Natal (13)**

Bergville - Dr. Ariena Shepherd

Bergville – Dr. Jubie Muller

Camperdown – Dr. Anthony van Tonder

Dundee – Drs. Marais, Fynn and Reynolds

Eshowe – Drs. Pryke and Hoffman

Estcourt – Drs. Turner, Tedder, Taylor, Tratschler, Van Rooyen and Alwar

Kokstad – Drs. Clowes and Shrives

Mooi River – Drs. Fowler, Hartley, Alexander and Reisinger

Mtubatuba – Dr. Trever Viljoen

Pietermaritzburg – Dr. Phillip Kretzmann

Pongola – Dr. Heinz Kohrs

Underberg - Drs. Collins, King and Delaney

Vryheid – Drs. Theron and Theron

**Eastern Cape (16)**

Alexandria - Dr. Johan Olivier

Alexandria – Dr. Charlene Boy

Aliwal North – Drs. Troskie and Strauss

Bathurst – Dr. Jane Pistorius

Cofimvaba – Dr. Werner Wentzel

Cradock – Dr. Frans Erasmus

Graaff-Reinet - Dr. Roland Larson

Humansdorp – Drs. Van Niekerk, Jansen Van Vuuren and Davis

Jeffreys Bay – Drs. Lategan, Hoek and McFarlane

Middelburg/Steynsburg – Drs. Van Rooyen and Viljoen

Port Alfred – Dr. Leon de Bruyn

Queenstown – Drs. Du Preez, Godley, Klopper, Jansen van Vuuren, De Klerk and Catherine

Somerset East – Drs. Farrel, Louw and Ross

Stutterheim – Dr. Dave Waterman

Uitenhage – Drs. Mulder and Krüger

Witelsbos – Dr. Elmien Kotze

**Western Cape (23)**

Beaufort West - Dr. Jaco Pienaar

Beaufort West – Dr. Bennie Grobler

Caledon – Drs. Retief, Coetzer and Janssen

Caledon – Drs. Louw and Viljoen

Ceres – Drs. Pieterse, Wium, De Villiers and Scheepers

Darling – Drs. Van der Merwe, Adam and Senekal

George – Drs. Strydom, Truter and Pettifer

Heidelberg – Dr. Albert van Zyl

Malmesbury – Dr. Otto Kriek  
Malmesbury – Dr. Markus Fourie  
Malmesbury – Dr. Andries Lesch  
Malmesbury – Drs. Heyns and Zolner  
Oudtshoorn – Dr. Glen Carlisle  
Oudtshoorn – Dr. Adriaan Olivier  
Piketberg – Dr. André van der Merwe  
Plettenberg Bay – Dr. André Reitz  
Riversdale – Drs. Du Plessis, Taylor and De Bruyn  
Stellenbosch – Dr. Alfred Kidd  
Swellendam – Dr. Jacques Malan  
Tulbagh/Ceres – Drs. Hamman, Wilson and Triegaardt  
Vredenburg – Dr. Izak Rust  
Wellington – Dr. Van Zyl and Louw  
Worcester- Dr. Kobus Rabe

#### **Northern Cape (7)**

Calvinia – Dr. Bertus Nel  
Colesberg – Drs. Rous and Rous  
De Aar – Dr. Donald Anderson  
Kathu – Dr. Jan Vorster  
Kimberley – Drs. Van Heerden and Swart  
Postmasburg – Dr. Boeta van der Merwe  
Upington – Drs. Vorster and Visser

#### **Feedlots (1)**

Drs. Morris and Du Preez

#### **Laboratory reports (5)**

Dr. Marijke Henton - Vetdiagnostix, Johannesburg  
Dr. Liza du Plessis – Idexx SA - Johannesburg  
Dr. Last – Vetdiagnostix, Pietermaritzburg  
Dr. Sophette Gers – Pathcare, Cape Town  
Queenstown Provincial Laboratory

## **Summary**

**WORLD ANIMAL WELFARE DAY**

October 6<sup>th</sup> is World Animal Welfare day. Whether it's livestock or other animals, all of us who deal with animals should ask ourselves some questions. What am I doing to maintain or improve the welfare of animals? Could I do more? Am I active enough in encouraging and persuading others to care better for their animals? Where are the biggest problems, and how can they be overcome?

Livestock Welfare, production and profitability are largely interlinked and so looking after welfare does not come at the price of a good farm management, in fact looking after welfare also benefits production and profits.

We at Livestock Welfare Coordinating Committee would like to know what you (or your organisation) do to promote the compassionate and caring management of livestock under your control – please email us at [secretary@lwcc.org.za](mailto:secretary@lwcc.org.za).

Gareth Bath  
Professor Emeritus  
Chair: LWCC

### **IMPORTANT MESSAGE**

**Animal welfare becomes more and more important!**

**Visit the website of the Livestock Welfare Coordinating Committee  
([www.lwcc.co.za](http://www.lwcc.co.za))**

### **Animal Identification, Tracibility and Disease control**

I recently attended the Red Meat Producers Organization's national congress and the ALPHA EXPO in Parys. Common goals expressed are that we should get our house in order so that producers can export meat.

Trading partners need to know that South Africa has a system in place where animals are identified on a central data base and movement can be traced. A public private process is on its way putting this system in place.

All of us (Farmers, State and Private Veterinarians) should all work together to make sure that diseases such as Foot and Mouth Disease and other controlled and notifiable diseases are eradicated and/or under control!

Visit the veterinary strategy for action plans

<http://nahf.co.za/wp-content/uploads/Vet-strategy-final-signed.pdf>

<http://nahf.co.za/controlled-and-notifiable-diseases/>

It is in the interest of all cattle farmers to ensure that herds (bulls) are free of trichomonosis! This is an area disease.

Breeding societies should get Vendor's declarations in place and buyers should demand declarations when buying animals!

When buying animals this Vendor declaration can help you to minimize risk!

#### **VENDOR DECLARATION BOVINE BRUCELLOSIS**

I hereby declare that I am the legal owner or authorised representative of the cattle on sale and am competent to make this declaration

1	The cattle for sale are clearly and permanently identified		Yes	No
2	The cattle for sale/slaughter were born on my farm		Yes	No
3	The farm has a closed herd policy i.e. I do not buy in cattle, rent out grazing or speculate with cattle		Yes	No
4	I practice bio-security on my farm to a level that is **	Poor	Moderate	Good
5	I vaccinate my heifer calves against Bovine Brucellosis once between the ages of 4 – 8 months		Yes	No
6	In addition I vaccinate my cattle older than 8 months with RB51		Yes	No
7	I have all the cattle on my farm tested for Bovine Brucellosis		Yes (date)	No
8	My herd has been tested negative within the past year		Yes	No
9	I did not buy in cattle since my last negative brucellosis test		Yes	No

10	I/my vet investigates any abortions on my farm		Yes	No
11	To the best of my knowledge, my immediate neighbours and farms in my area are free of Bovine Brucellosis		Yes	No
12	I use a veterinarian to advise me on my cattle's herd health		Yes	No
13	The cattle handling facilities on my farm are	Poor	Average	Good

Note: Vaccination does not mean freedom from Bovine Brucellosis as cattle can still be carriers  
Please attach the most recent *Brucella* blood test certificate

Owner or authorised representative:.....

Signature:.....

Date:.....

**\*\* \* Biosecurity**

Poor – speculates with cattle, does not vaccinate, poor fences, cattle come into contact with other cattle

Medium – Vaccinates heifers, does not buy in cattle of unknown health status

Good – closed herd/never buys in cattle, vaccinates heifers and no contact with other cattle, follows a herd health plan as advised by his veterinarian, does not allow transport trucks onto property, washes and disinfects truck after returning from the abattoir or auction grounds.

Compiled by: Dr. Sewellyn Davey, Chairman of the Brucellosis Steering committee of the National Animal Health Forum

## **OVINE JOHNIE'S DISEASE VENDOR DECLARATION**

### **ON THE SALE OF SHEEP**

(Updated Draft May 2015)

- |   |  |     |    |
|---|--|-----|----|
| 1. I hereby declare that I am the owner or authorised representative of the sheep on sale and am competent to make this declaration.                    | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 2. The sheep for sale are clearly identified in the accompanying description.   | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 3. The sheep for sale were born on my farm.   | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 4. The farm has a closed flock policy. (No live sheep are brought onto the farm from elsewhere)   | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 5. I know the signs of the disease and to the best of my knowledge, all of my properties are free of cases of Ovine Johnie's Disease.                   | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 6. I have actively looked for Ovine Johnie's Disease and have had tests done for this.  | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |
| 7. To the best of my knowledge, my immediate neighbours and farms in my magisterial district of my farm(s) are free of cases of Ovine Johnie's Disease. | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO |
| YES   | NO   |     |    |

- |   |  |     |       |
|---|--|-----|-------|
| 8. The sheep on my properties have been vaccinated against Ovine Johnie's Disease and are clearly marked with the approved ear tag. | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO    |
| YES   | NO   |     |       |
| 9. All lambs born are vaccinated  | <table border="1"><tr><td>YES</td><td>NO</td></tr></table> | YES | NO    |
| YES   | NO   |     |       |
| 10. If vaccinated, the number of years that the vaccinations have been done is  | <table border="1"><tr><td></td><td>years</td></tr></table> |     | years |
|   | years  |     |       |

**NOTE:** Vaccination does not mean freedom from OJD, vaccinated animals can still be carriers.

Statement 8 and 9 apply only to already infected flocks, and such sheep can only be sold to other infected flocks by law.

Buyers should consult their veterinary advisor before any purchases.

Signature \_\_\_\_\_

Date \_\_\_\_\_

NAME \_\_\_\_\_

Farm: \_\_\_\_\_

District: \_\_\_\_\_

OWNER OR AUTHORIZED  
REPRESENTATIVE

The use of this declaration is supported by the following organisations:



UNIVERSITEIT VAN PRETORIA  
UNIVERSITY OF PRETORIA  
UNIVERSITEIT VAN PRETORIA



**RUVASA**  
Rural Veterinary Association of South Africa





# **In May there had been an outbreak of Rift Valley Fever in the Jacobsdal area of the Free State. Be prepared for the next outbreak in your area!!**

Rift Valley Fever is a zoonotic disease.

“The symptoms are suggestive of influenza, have a rapid onset and consist of rigors, fever, headache and pain in muscles, back and joints. Many people recover uneventfully but a variable percentage may develop more severe symptoms, the liver may be affected and the patient is inclined to haemorrhage. Several other symptoms may also follow infection, including involvement of the eyes.”  
(J H du Preez, et al, 2017)

Farmers are advised to vaccinate their animals against Rift Valley Fever. Live vaccines can only be used in non-pregnant animals as the live vaccine can cause abortions.

Only dead inactivated vaccines must be used in pregnant animals.

Please take care when vaccines are administered after outbreaks have been detected. Use a sterile needle for each animal as the virus can be spread between animals if the animals are in the incubation period. Animals in the incubation period will have a virus infection but will not show symptoms of the disease yet.

## **Bovine brucellosis control should continuously be on the mind of every cattle and game farmer!**

Get involved in controlling Bovine Brucellosis – visit [www.nahf](http://www.nahf), click on Information centre, click on Brucellosis.

### **Why should I test my cattle for brucellosis?**

- Brucellosis is a herd disease – this means if you even have a single infected animal, it affects the status of your whole herd! An infected cow will rapidly infect the rest of your herd if not identified and slaughtered before calving.
- Brucellosis causes reproduction and production losses in affected herds, which decreases financial turnover.
- Brucellosis is a misleading disease as abortions may not always occur, abortions are not always noticed and affected cattle may appear visibly healthy. Only through testing can brucellosis be confirmed.

- The disease is a zoonosis which can infect you, your family and your workers. Infection may lead to chronic and debilitating disease.
- If you have recently bought in animals from an auction or a herd with no proof that the herd of origin was recently tested for bovine brucellosis (CA3 declaration), you could have bought in positive or latently positive animal(s).
- If neighbouring farms or farms you access for grazing are positive for brucellosis, your cattle could have contracted the disease.
- If you do not know your status and you are farming with brucellosis unknowingly, you are contributing to the brucellosis risk of our national cattle herd.

Always be aware of all possible ways how your herd can be become infected with the bacteria, *Brucella abortus*.

- **Just one miss step may be the beginning of misery!**
- **Remember this is a herd disease!**
- **If one animal is tested positive, the herd is seen as positive until proved to be negative according to the brucellosis scheme!**
- **Brucellosis is a State controlled disease.**
- **Brucellosis is a zoonosis – a disease transmitted from animals to humans**

**A check list has been drawn up to help you calculate what the risk is of bringing the *Brucella* bacteria onto your farm.**

Below is a map of new outbreaks of bovine brucellosis as supplied by DAFF.

It should be emphasized that all cases of State controlled and notifiable diseases have to be reported to the State Veterinary services!

<http://nahf.co.za/controlled-and-notifiable-diseases/>

It is with great concern that animals that have aborted are sold at auctions without the herd and specific animals are tested for bovine brucellosis! Brucellosis is a herd disease.



8	Data base of all animals		
9	Closed herd		
10	When last were animals bought in or moved from another farm?		
11	Only buy in animals from a farm which has a recent negative tested brucellosis herd certificate		
12	Origin(s) of acquired cattle? Bought at an auction?		
13	Keep heifers separate from herd until they have calved and tested negative for brucellosis		
14	Quarantine camp available		
15	Separate calving camps		
16	Were all heifers vaccinated between 4 and 8 months vaccinated with Strain 19 or RB51?		
17	Any cattle vaccinated with Strain 19 over 8 months of age? History over last few years.		
18	Were there any abortions on the farm – samples taken, diagnosis?		
19	All sexually mature cattle in herd tested for bovine brucellosis (provide proof)		
20	Bovine brucellosis is a State controlled disease. Positive cattle are branded with a C on the right side of the neck.		
21	Isolation of infected animals & separate handling facilities		
22	Prohibition of movement of animals off quarantined property except under cover of a Red cross permit for slaughter at an abattoir		
23	Prohibition of use and on-farm disposal of unboiled, unpasteurised or unsterilised milk on quarantined property		
24	Disinfection of places where infection is a possibility.		
25	Neighbours/ recent buyers informed of infected herd status		
26	Fly, crow and predator control		
27	Destruction of afterbirths/abortions in a responsible manner		
28	Beware of livestock, game interface		

**Below are short and longer versions on using vaccines to help control brucellosis. Please spread this information as wide as possible.**

## **Brucellosis**

### **Play your part in the war on brucellosis – don't wait until it's too late; vaccinate!**

Bovine brucellosis, caused by *Brucella abortus*, is reported across all 9 provinces of South Africa especially in the central and Highveld regions. Bovine Brucellosis mainly causes abortion in cattle and can infect most other mammals, including humans.

Brucellosis may occur in persons who are exposed to infected animals, particularly through aborted and normal fetal material of infected cows, and through the consumption of unpasteurised milk. Symptoms in infected persons include profuse sweating mostly during the night, fever, extreme tiredness, aches in bones and joints, especially the lower back, hip or knee joints.

All female cattle between the ages of 4-8 months have to be vaccinated against brucellosis with a registered product to help protect the national cattle herd.\* Vaccination helps to decrease shedding of *Brucella* bacteria from infected animals – this helps to limit the spread of brucellosis within a herd and decreases the infection pressure. Always use vaccines according to the manufacturer's instructions. Contact your private veterinarian or state veterinary services for more information.

#### **5 keys to controlling brucellosis:**

- Vaccination is one of the important tools used to control and prevent brucellosis. Vaccination of all heifers between 4-8 months is required. Do not vaccinate male cattle.
- Know your status - test your cattle herd for brucellosis.
- Only buy cattle from recently tested, brucellosis negative herds.
- Don't share grazing with untested cattle and ensure that your fences are intact.
- If you have brucellosis in your herd, follow the advice of your state veterinarian to get rid of the infection as quickly as possible (branding, separation, vaccination and slaughter). Delays in action against the disease will increase losses and prolong quarantine.

Let's all play our part to protect each other – our cattle's health, our own health and our livelihoods.

\*Brucellosis is a controlled disease under the Animal Diseases Act, 1984 (Act No 35 of 1984).

### **Play your part in the war on brucellosis**

- Don't wait until it's too late - vaccinate! To protect the national cattle herd, all female cattle between the ages of 4-8 months have to be vaccinated against brucellosis with a registered product.\*
- Vaccination strengthens immunity against brucellosis and decreases bacterial shedding.
- S19 vaccine for cattle (Reg. No. G0101. Act 36/1947) or RB51 (Reg. No. G 3056. Act 36/1947) can be used. Always use vaccines according to the manufacturer's instructions. Contact your private veterinarian or local state veterinary services for more information.

- Know your herd's status by testing. Know the status of herds that you buy or receive cattle from by requesting herd test results.
- Let's all play our part to protect each other – our cattle's health, our own health and our livelihoods.

\*Brucellosis is a controlled disease under the Animal Diseases Act, 1984 (Act No 35 of 1984).

At the following link information needed to educate cattle owners and consumers of unpasteurised dairy products regarding brucellosis are available..

<http://nahf.co.za/category/diseases/brucellosis/>

Translations in the major languages regarding the 5 core facts on bovine brucellosis are available at the following link:

<http://nahf.co.za/translations-for-bovine-brucellosis-5-core-facts-and-a-dozen-things/>

Websites that are there to help you with information regarding animal health:

## National Animal Health Forum

[www.nahf.co.za](http://www.nahf.co.za)

Read what the Forum is all about:

<http://nahf.co.za/about/>

This website will become the information centre of animal health in Southern Africa.

On the toolbar click on **Stakeholders** and you will find links to producer organizations and other organizations who are participating in the NAHF

<http://nahf.co.za/stakeholders/>

Provincial Animal Health Forums have their own site – click on **Provinces** <http://nahf.co.za/provinces/>

Important is to study the Veterinary Strategy (2016 -2026) as it gives direction to where we are going with Animal Health in South Africa.

<http://nahf.co.za/wp-content/uploads/Vet-strategy-final-signed.pdf>

Click on **Info centre** for more information on the “war” we have against Bovine Brucellosis. Please be up to date on the role all have to play to control this zoonotic disease.

<http://nahf.co.za/category/diseases/brucellosis/>

Information on other controlled diseases (Ovine Johne's Disease, Pest of small stock – PPR, and African Horse Sickness) is available.

This link will continuously be updated.

Information on **antibiotic resistance** is also available at this address:

<http://nahf.co.za/category/antibiotic-resistance/>

## Rural Veterinary Association of South Africa

[www.ruvasa.co.za](http://www.ruvasa.co.za)

Click on **Disease reporting** where maps and information can be sourced on the prevalence of diseases in all provinces. Abattoir reports are available. Use the information available to update management programmes

## Landbouweekblad's webpage

[www.landbou.com](http://www.landbou.com)

[Vra vir Faffa](#)

Click on: **Indeks van antwoorde** where more than 4 000 answers can be sourced on animal health.

Click on Beeste

Click on Siektes

Click on Brusellose

Stop Brusellose

Gevaar om Beesbrusellose (BBR) deur vendusies en skoue te versprei

Rapportering aan bure of ander eienaars oor die voorkoms van brusellose

Inligting oor brusellose op die NAHF se webblad

Kuddebestuur voor die dekseisoen

Bees Brusellose handleiding

Teenliggaamwaardes om beesbrusellose in koeie te bepaal

Veterinêre Strategie 2016 -2026

'n Dosyn dinge wat jy moet weet van beesbrusellose

Vyf kernfeite wat jy van beesbrusellose (Besmetlike misgeboorte – BM) behoort te weet

Veiligheid van vleis en biltong afkomstig van 'n bees met brusellose

Vervoer van diere uit 'n positiewe brusellose kudde

Beheer van brusellose in 'n beeskudde

Boerderypraktyke wat die gevaar van die voorkoms van brusellose verhoog

Pak brusellose by die horings

Brucellose kan jou lewe verwoes

Brusellose in wild

Bestuur van positiewe besmetlike misgeboorte beeste

Aankoop van beeste wat besmetlike misgeboorte het

Antwoorde oor brusellose

Behandeling van besmetlike misgeboorte

Besmetlike misgeboorte uitbreek in 'n kudde

Gevaar van brusellose onderskat

RB51-inenting teen brusellose in dragtige koeie

Alles oor Besmetlike Misgeboorte (BM)

Kompensasie vir BM en TB positiewe beeste?

Nóg vrae oor besmetlike misgeboorte

Koeie positief getoets vir besmetlike misgeboorte

Vrae, antwoorde oor besmetlike misgeboorte

Brucellose: Wat staan ons te doen?

## Internal parasite control

[www.wormx.info](http://www.wormx.info)

## Summary of disease report for August 2018

120 Reports from veterinary practices and laboratories were received (Mpumalanga (MP) 9; Gauteng (G) 7; Limpopo (L) 9; Northwest (NW) 10; Free State (FS) 20; KwaZulu-Natal (KZN) 13; Eastern Cape (EC) 16; Western Cape (WC) 23; Northern Cape (NC) 7; Feedlots (FL) 1 and Laboratories (Lab) 5).

For the detailed report and previous reports go to [www.ruvasa.co.za](http://www.ruvasa.co.za) and click on Disease reporting

## Internal parasites

The following reports were received from practices regarding internal parasite infestations:

Internal parasites	MP	G	L	NW	FS	KZN	EC	WC	NC
Roundworms	x	x		x	x	x	x	x	
Resistant roundworms	x	x							
Wireworm		x		x	x	x	x	x	x



Brown stomach-worm								X	
Long-necked bankruptworm									
Large-mouthed bowelworm									
Nodularworm							X		
Lungworm									
Eyeworm									
<i>Parafilaria</i>						X			
Tapeworms	X	X			X				
Liver fluke	X			X	X	X		X	
Conical fluke	X			X	X	X			
Cysticercosis (measles)	X		X	X					X
Schistosomiasis (bilharzia)									
Coccidiosis	X	X		X	X		X	X	X
Cryptosporidiosis				X	X	X	X	X	

Good rainfall occurred in many areas of the the winter rain fall area. Be aware of brown stomach worm infestation that may become a problem after the severe drought. Use the five point check to keep on top of what is happening in the flock. For further detail contact your local veterinarian.

<http://hulp.landbou.com/kundiges/vra-vir-faffa/vyfpuntplan-en-famacha-stelsel-vir-inwendige-parasietbestuur-in-skape/>

[https://docs.wixstatic.com/ugd/ad98\\_cb447e77eef6450f93a2b23cb0e6b9de.pdf](https://docs.wixstatic.com/ugd/ad98_cb447e77eef6450f93a2b23cb0e6b9de.pdf)

Wireworm infestations have been reported from 7 provinces. Most farmers think that wireworm is dormant in winter. Check animals for bottle jaw and anaemia. As temperatures rise, be aware of early outbreaks of wireworm infestations. For worm eggs to hatch humidity, temperature above 15 degree Celsius and oxygen are needed.

It cannot be stressed enough that the basis of preventing cryptosporidiosis and pathogenic *E. coli* outbreaks is biosecurity and making sure that the monthers give the best possible colostrum quality to their offspring.

Cryptosporidiosis in combination with pathogenic *E. coli* is still causing huge problems for cattle, sheep and goat farmers. Young animals are extremely vulnerable. Biosecurity measures should be kept at a high level at all times. The immunity of ewes and cows should be kept at the highest level so that colostrum (passive immunity) could protect their offspring.

It is important to contact your veterinarian to help you diagnose what the problem(s) on the farm is when young animals start dying. Samples are set away for diagnosis. *Cryptosporidia* and *E.coli* can be present in the water and food sources. Supply workers with protective clothing which have to be kept on the farm.

Choose disinfectants that is registered for killing *Cryptosporidium*!

## External parasites

The following reports were received from practices regarding external parasite infestations:

External parasites	MP	G	L	NW	FS	KZN	EC	WC	NC
Blue ticks	x	x		x	x	x	x	x	
Resistant blue ticks								x	
Heartwater ticks	x	x		x		x	x		
Brown ear-ticks		x				x			
Bont-legged ticks	x			x	x				x
Red-legged ticks	x			x	x				
Paralysis ticks					x		x		
Tampans									
Biting lice				x	x	x		x	
Sucking lice				x	x	x	x		
Itch mites									
Sheep scab		x		x			x	x	
Mange mites	x	x		x	x	x	x	x	

Nuisance flies	x					x		x	
Midges						x		x	
Mosquitoes									
Blowflies			x						
Screw-worm	x							x	
Gedoelestia (uitpeuloogsiekte)									
Nasal bot	x				x				x

Blue tick infestations were reported from all provinces. Blue ticks (African and Asiatic blue ticks) are able to transmit red water, anaplasmosis and lumpy skin disease.

Make sure to assess the blue tick resistance status on your farm before buying tickicides. Your veterinarian will be able to collect engorged blue ticks to be tested for resistance.

Actives to be tested for resistance are: organophosphates, pyrethroids, amidines, fipronil. Actives registered only for controlling blue ticks are: macrocyclic lactones, fluazuron (acaricide growth regulator).

Discuss your tick control programme with your veterinarian as controlling ticks early in spring can prevent large outbreaks of ticks in the summer.

Below is a list of diseases transmitted by ticks.

## Tick borne diseases

The following tick-borne diseases were reported by practices in the provinces:

Tick borne diseases	MP	G	L	NW	FS	KZN	EC	WC	NC
African red water	x		x	x	x	x	x	x	
Asiatic red water	x	x	x		x	x	x	x	
Anaplasmosis		x		x	x	x	x	x	
Heartwater	x	x	x	x		x	x		
Lumpy skin disease				x	x	x	x		

Corridor disease									
Theileriosis									

Asiatic red water is spreading and is one of the deadliest diseases in cattle.

Numerous mortalities were reported!

The keyword is **vaccinate** your animals! Contact your veterinarian.

Anaplasmosis outbreaks were reported in 6 provinces.

The following tick toxicosis was reported by practices in the provinces:

Tick toxicosis	MP	G	L	NW	FS	KZN	EC	WC	NC
Sweating sickness				x	x			x	

## Insect transmittable diseases

The following insect transmittable diseases were reported by practices in the provinces:

Insect transmittable diseases	MP	G	L	NW	FS	KZN	EC	WC	NC
Lumpy skin disease				x	x	x	x		
Ephemeral fever (Three day stiff sickness)						x			
Blue tongue	x	x			x				
Rift Valley Fever									
Wesselsbron									
Nagana						x			

Insect borne diseases have decreased dramatically after colder weather occurred in most parts of the country. Now is the time to vaccinate your animals against these diseases.

**Have you vaccinated your animals vaccinated against Rift Valley Fever?**

**Plan now, with the assistance of your veterinarian, the vaccination programme during the winter months, to prevent outbreaks of diseases during the next rainy season.**

**Out of experience I can tell you that when outbreaks of diseases occur, vaccines will be difficult to acquire as many people will be ordering vaccines all at once.**

## **Venerial diseases**

The following venereal diseases were reported by practices in the provinces:

<b>Venereal diseases</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Trichomonosis</b>	<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>		<b>x</b>
<b>Vibriosis</b>			<b>x</b>	<b>x</b>	<b>x</b>		<b>x</b>	<b>x</b>	
<b>Pizzle disease</b>							<b>x</b>	<b>x</b>	<b>x</b>
<b><i>Actinobacillus seminis</i></b>				<b>x</b>					

New cases of **trichomonosis** are reported every month and this disease is out of control. Make sure to buy bulls from farmers where biosecurity measures are in place and bulls are tested for these diseases at regular intervals.

Make sure that fences are in tact and gates closed so that bulls cannot escape to neighbouring cows that may be infected with *Trichomonas* and become infected or infected neighbouring bulls are jumping fences.

Cattle study groups should discuss preventative and control measures with their veterinarians. **Be sure to test bulls regularly for these diseases.**

**Beware when buying in or sharing bulls! Remember female animals may also be infected.**

Study the Good management SOP's for cattle farmers on the RPO website

<http://www.rpo.co.za/wp-content/uploads/2016/04/nuutRPO-NERPO-Code-Addendum.pdf>

<http://www.rpo.co.za/wp-content/uploads/2016/04/nuutRPO-NERPO-Code-Addendum-4-Good-management-practices-and-SOPs-for-cattle-farmers-1.pdf>

Consider Trichomonosis as an area diseases, farmers should work together to keep areas free from diseases such as trichomonosis, brucellosis and sheep scab.

## **Bacterial diseases**

The following bacterial diseases were reported by practices in the provinces:

<b>Bacterial diseases</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Anthrax</b>									
<b>Blackquarter</b>	<b>x</b>			<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>		
<b>Botulism</b>					<b>x</b>	<b>x</b>	<b>x</b>		
<b>Pulpy kidney</b>		<b>x</b>			<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	
<b>Lamb dysentery</b>							<b>x</b>		<b>x</b>
<b>Swelled head</b>		<b>x</b>						<b>x</b>	
<b>Red gut (cattle)</b>		<b>x</b>			<b>x</b>	<b>x</b>		<b>x</b>	
<b>Blood gut (sheep)</b>	<b>x</b>			<b>x</b>	<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>
<b>Tetanus</b>		<b>x</b>						<b>x</b>	
<b>Salmonellosis</b>	<b>x</b>		<b>x</b>	<b>x</b>					
<b>Bovine brucellosis</b>				<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	
<b>Ovine brucellosis (Ram's disease)</b>					<b>x</b>			<b>x</b>	
<b>Bovine tuberculosis</b>								<b>x</b>	
<b>Johne's</b>								<b>x</b>	
<b>Leptospirosis</b>									
<b>Listeriosis</b>									
<i>Pseudomonas</i>									
<i>Fusibacterium necrophorum</i>								<b>x</b>	
<b>Septicaemia</b>		<b>x</b>				<b>x</b>			<b>x</b>
<i>E. coli</i>	<b>x</b>	<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>
<b>Enzootic abortion</b>	<b>x</b>		<b>x</b>		<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>
<b>Lumpy wool</b>					<b>x</b>				

Uterine gangrene							x	x	x
Bovine dermatophilosis (Senkobo disease)						x			
Wooden tongue									
Lumpy jaw									

Multiclostridial vaccines should be used if blackquarter outbreaks still occur when only using a vaccine containing *Clostridium chauvoei*. Remember to give a booster vaccine when using an inactivate vaccine for the first time. **Read the packet insert!!** Study the table above and determine the risk for animals on your farm. Get advice from your veterinarian on *Cryptosporidium*/*E. coli* outbreaks in your area and what to do to prevent losses in lambs and calves.

## Viral diseases

The following viral diseases were reported by practices in the provinces:

Viral diseases	MP	G	L	NW	FS	KZN	EC	WC	NC
BMC (snotsiekte)	x		x	x	x	x	x	x	
Rabies (cattle)				x			x		
BVD			x	x				x	x
IBR							x	x	x
BRSV									
PI3									
Maedi visna virus									
Rotavirus / Coronavirus						x		x	
Enzootic bovine leucosis (EBL)				x			x	x	
Sheep leucosis									
Jaagsiekte		x							

<b>Orf</b>	<b>x</b>	<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>
<b>Warts</b>				<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>

There is no treatment for viral diseases with the result that animals have to be protected by vaccinations if they are available.

Preventative vaccinations are the best way to protect animals against viruses and bacteria causing pneumonia.

Keep cattle and wildebeest well separated especially when wildebeest are under stress!

Discuss vaccination programmes and biosecurity measures with your veterinarian.

## Fungal diseases

The following fungal disease was reported by practices in the provinces:

<b>Fungal diseases</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Ringworm</b>	<b>x</b>	<b>x</b>		<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	<b>x</b>	

## Protozoal diseases

<b>Protozoal diseases</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Besnoitiosis (olifantsvelsiekte)</b>		<b>x</b>							

## Toxicities

The following toxicities were reported by practices in the provinces:

<b>Toxicities</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Cardiac glycoside</b>							<b>x</b>	<b>x</b>	
<b>Slangkop</b>									
<b>Crotolaria</b>									
<b>Gifblaar</b>		<b>x</b>							









**Beware when buying in animals or moving them into rested grazing camps as they are the animals which usually eat toxic plants such as tulp and ink berries (*Cestrum*).**

**Do have activated charcoal on the farm as the antidote for tulip poisoning! **Seven provinces reported tulip toxicity!****

**Toxic plants are sometimes eaten by young animals that do not know these plants. Be aware of this situation and know where these plants are growing on the farm.**

**For further information on treatment of tulp and other toxicities visit:**

**[www.landbou.com](http://www.landbou.com)**

**Vra vir Faffa**

**Klik op Indeks van antwoorde**

**Klik op Beeste of Skape**

**Klik op Vergiftigings**

**Klik op die Opskrifte**

**Every month there are reports of urea poisoning. Be aware when feeding this product that the correct concentration is used and that the lick does not get wet!**

**A few cases of Lantana-poisoning were reported.**

**Research are being done to control Lantana:**

We would like to investigate involvement of your readers and yourself in the development of a National Programme for Management of Lantana similar to the attached National Programme for Management of Parthenium. Our focus would be on the biological control of the species, however, farmers, landowners and communities would be interested in an integrated approach to the management of the species. Please do bear in mind that the rust-fungus will unfortunately not be a 'silver-bullet' as it is likely to impact some subspecies more than others and work better in some micro-climates than others..

Please can we consider how your readers would be able to contribute to the development of a National Programme? One element would be accurate mapping of the distribution of Lantana. If readers could be encouraged to report locations of Lantana then a more comprehensive map of its distribution would be feasible (we need to make sure that this is done in a co-ordinated fashion and using technology that allows for accuracy and ease of data collection (smart phone application to geographically referenced database – which would need to be set up and managed).

Encouraging readers to give input into a National Programme would result in greater support for its implementation. We would need to make sure that this is not too tedious a process.

[illegible]

<b>Selenium</b>				<b>X</b>		<b>X</b>	<b>X</b>		
<b>Magnesium</b>				<b>X</b>	<b>X</b>				
<b>Manganese</b>						<b>X</b>			
<b>Vitamin A</b>				<b>X</b>	<b>X</b>	<b>X</b>		<b>X</b>	
<b>Vitamin B 1</b>								<b>X</b>	

There are antagonists such as calcium, iron and sulphur which hamper the uptake of micro-minerals. Have water and soil samples analysed to see what the levels of these antagonists are. Arrange with your veterinarian to have liver samples analysed to determine the status of these micro-minerals in your herd or flock.

Selenium is a powerful anti-oxidant and necessary for immunity. Check the status of the herd.

Beware of fluoride poisoning as borehole water levels drop.

Supplement animals with vitamin A during winter and drought conditions.

## Multifactorial diseases and other conditions

The following conditions were reported by practices in the provinces

<b>Multifactorial diseases and other conditions</b>	<b>MP</b>	<b>G</b>	<b>L</b>	<b>NW</b>	<b>FS</b>	<b>KZN</b>	<b>EC</b>	<b>WC</b>	<b>NC</b>
<b>Abortions</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Stillbirths</b>		<b>X</b>		<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	
<b>Abscesses</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Intestinal ulcers</b>				<b>X</b>				<b>X</b>	
<b>Bladder stones –urolithiasis</b>									
<b>Blindness</b>					<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Bloat</b>					<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>
<b>Blue udder</b>					<b>X</b>		<b>X</b>	<b>X</b>	<b>X</b>
<b>Diarrhoea</b>	<b>X</b>			<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	<b>X</b>	
<b>Epididymitis</b>	<b>X</b>	<b>X</b>		<b>X</b>	<b>X</b>			<b>X</b>	

Eye cancer		x			x		x	x	
Eye infections		x		x	x	x	x	x	x
Joint ill	x				x	x	x	x	
Lameness/foot problems	x	x		x	x	x	x	x	
Lung infection	x	x	x	x	x	x	x	x	x
Mastitis	x	x		x	x	x	x	x	
Navel ill	x	x				x	x	x	
Umbilical hernia		x							
Red gut (sheep, torsion of gut)								x	
Rectal prolaps		x							
Swelsiekte									
Traumatic reticulo-pericarditis							x	x	
Trauma						x			x
Teeth wear									
Plastic bags (ingestion)									
Downer	x	x			x	x	x	x	
Anaphylactic shock									
Vestibular syndrome (middle ear infection)									

Discuss the origin, treatment and prevention of these diseases with your veterinarian.

The cause of abortions should be established: brucellosis, enzootic abortion, Q-fever, leptospirosis, etc. The necessary preventative measures can then be taken.

Pneumonia and lameness (foot conditions) are wide spread.

## Metabolic diseases

The following diseases were reported by practices in the provinces:

Metabolic diseases	MP	G	L	NW	FS	KZN	EC	WC	NC
Acidosis	x	x		x	x	x		x	
Displaced abomasum					x		x	x	
Ketosis (Domsiekte)		x			x		x	x	x
Milk fever		x			x	x	x	x	

Make sure that you adapt animals to feed containing concentrates as more and more cases of acidosis are reported when grazing animals on harvested maize fields.

Discuss the etiology, treatment and prevention of these diseases with your veterinarian.

## Reproductive diseases

Reproductive diseases	MP	G	L	NW	FS	KZN	EC	WC	NC
Dystocia (difficult births)	x	x	x	x	x	x	x	x	x
Endometritis		x			x		x	x	
Hydrops									
Metritis	x			x	x	x	x	x	
Poor conception	x			x	x	x		x	x
Retained afterbirth	x	x		x	x	x	x	x	x
Sheath prolaps		x		x	x	x	x		x
Uterine prolaps	x	x	x	x	x	x	x	x	x
Vaginal prolaps	x	x		x	x	x	x	x	
Penis injury									
Orchitis									

A poor conception rate on many farms is a huge issue. Visit your veterinarian to rectify this problem.

## Environmental conditions



	MP	G	L	NW	FS	KZN	EC	WC	NC
Exposure to cold					x		x	x	
Frozen to death					x				
Heat stress									
Lightning									
Electrocution									
Drought							x		

Be prepared when cold spells and snow fall are forecasted.

## Other conditions

	MP	G	L	NW	FS	KZN	EC	WC	NC
Drug residues (milk, meat, liver, kidney etc)									
Predators	x				x		x		
Theft			x		x	x			
Trauma (fractures etc)		x		x				x	
Trauma (veldfires)					x				

In the CODE OF CONDUCT of the RPO the following standard operating procedures are documented.

The local veterinarian should be your partner to help you achieve the necessary standards.

<http://www.rpo.co.za/BestPractices/English.aspx>

## PRECAUTIONARY MEASURES TO SUPPORT BIO-SECURITY.

Precautionary measures are required to protect the herd against diseases acquired because of external contact. The following categories are of concern:

### 1. DIRECT LIVESTOCK PURCHASES (and own animals returning):

The following should be **verified** before importing new animals into the herd:

How long animals have resided at the purchase or previous location?

Have there been any recent disease outbreaks in the location?

Do brand marks clearly confirm ownership?

Was a vaccination program followed (need paper or veterinarian proof). What are the local prevalent external parasites and the routinely implemented control program?

Is a veterinarian supported control program against transmittable diseases followed?

Dates and sufficient number of tests for reproductive diseases of both male and female

Dates and tests for zoonotic diseases

The above should also be verified with the purchaser's own veterinarian.

## **2. PURCHASES FROM SALES OR SPECULATORS**

Purchase only in areas which are not in close proximity to scheduled areas

Visually inspect the animals before purchasing for:

- \* brand marks

- \* parasite infestation

## **3. TRANSPORT TO THE FARM**

Use only reputable transporters

Has the truck been cleaned and disinfected?

Truck to follow the shortest uninterrupted route

Truck to take the shortest route to the handling facilities

Do not allow the truck personnel to get in contact with the farm herd

## **4. ARRIVAL ON THE FARM**

Off-load the livestock to limit stress and to be visually evaluated for any unnatural conditions.

Isolate them from the farm herd and shared facilities for at least 21 days (quarantine)

Retest for diseases of concern if needed, before mixing with the rest of the herd

Process new arrivals within 24 hrs after arrival (unique ID tag brand, dip, dose, vaccinate)

Inspect regularly

## **5. FEED PURCHASES**

Ensure bales of hay are sourced from areas that are not bordering scheduled areas

Purchase feed from reputable dealers only

Avoid buying feed in second hand bags

Ensure feed trucks are also disinfected and cleaned, especially if also used to transport animals to abattoirs

## **6. VISITORS**

Do not allow strangers or their vehicles amongst the livestock

Ensure fences are well maintained and preferably jackal and warthog proof

## **7. EMPLOYEES**

Do not allow the employees to eat in feed stores

Supply employees with sufficient ablution facilities

Regularly arrange to let employees be medicated for tape worm and have health check-ups

Keep record of all employee livestock on the property

Treat employee livestock with separate but dedicated health programs

Ensure employees understand the reason behind the implemented bio-security measures to help ensure compliance.

## **GENERAL AND REPRODUCTION MANAGEMENT**

Record keeping: All animals are individually identified and recorded.

To prove ownership: All animals are marked with the registered brand mark according to the Animal Identification Act, No 6 of 2002.

A defined breeding season is the basis of effective management: The breeding season coincides with the rainy season, i.e. the period when nutritive value of the pasture is at its best.

Sufficient energy reserves in the herd as measured by condition scoring are vital, especially for effective breeding, and when inadequate the herd is supplemented in consultation with a nutritionist: Condition scoring of bulls and cows are regularly done, particularly at the onset of the breeding season and supplemented if necessary.

Bull - cow ratios are maintained: A ratio of 1 to 25 is maintained in every separate herd.

Fertility of breeding bulls: All breeding bulls are tested for mating ability and semen quality before the breeding season.

Sexually transferable diseases: Sheath washes or scrapes on bulls are performed annually.

Diseases that can cause poor conception, abortion or weak calves: Cows are vaccinated against such diseases in consultation with the veterinarian.

Breeding success monitored by a veterinarian: Rectal pregnancy or scan diagnosis is done by the veterinarian 8 weeks after the breeding season.

Twenty percent of cows or more not pregnant: Further tests are done to determine cause of low pregnancy rate.

Culling of non-pregnant cows: Non-pregnant cows are removed from the herd and considered a necessary bonus to supporting herd income.

## **HERD HEALTH AND BIO-SECURITY**

Maintenance of herd health is key to a successful enterprise: A veterinarian should visit the farm bi-annually at least.

Calf mortality before 3 months of age is an important reason for poor weaning percentage: Good management practices are applied to limit early calf deaths.

Some diseases and parasites (internal and external) are more often encountered in specific areas: Annual vaccinations and a parasite control program should be applied according to regional requirements and in liaison with the veterinarian.

Farmers selling weaned calves to feedlots may want to have a market advantage compared to others: A specific vaccination program is applied before weaning for that purpose.

Herds may be at risk of being exposed to CA and TB: The herd is tested annually for CA and all heifers are vaccinated against CA between 4 and 8 months of age with an efficient, approved remedy. The herd is tested at least every 5 years for TB

Precautionary measures are required to prevent diseases being imported into the herd: A quarantine program to keep incoming animals separate is followed. All incoming animals have a suitable certificate of negative test results or are of a certified clean, closed herd.

Stock remedies and medicines should be registered, correctly stored and used before the transpire date:

All medicines and stock remedies are registered, stored and applied according to prescription.

Prescribed medicines with a specific application are under the control of the veterinary profession: All prescription medicines are obtained and applied under prescription from a veterinarian.

---

## **Practices that had nothing to report**

Cape Town – Sophette Gers

Kaapmuiden – Dr. Silke Pfitzer

Mokopane – Dr. Alwyn Venter (SV)

Nigel – dr. Cindy vd Westhuizen

Plettenberg Bay – Dr. André Reitz

Stutterheim – Dr. Dave Waterman

Vaalwater – Dr. Hampie van Staden

Vaalwater – Dr. Annemieke Müller

Vanderbijlpark – Dr. Kobus Kok

## **Ostriches**

### **Western Cape**

#### **Oudtshoorn**

Tapeworms – 2 Not a significant production problem but farmers like to Rx to get flocks clean. Most grower flocks rid themselves of infection

Red gut – 3 Overfeeding on maize and changes in intake behaviour. Overgrowth, Alpha toxin production. More adult birds. Treatment: tetracyclines effective.

Diarrhoea – 3 Associated with mouldy feed – drought, low cost and last stored roughage. Also cheaper maize results in feeding high volumes of maize, more of a acidosis condition underlying.

Yolk sack infection -3 Chick season is starting, open nest system with few drops of rain causes shell contamination. Weak bacteraemic chicks post hatch. Treatment effective with quinolones

Protein and energy deficiency - 3 Protein energy deficiency. Low feed intake due to feed differences (quality and source of roughage) short day periods for intake, high energy need during winter.

Mycotoxins – 2 DON high levels in feed. Last of season storage – desperate to obtain cheap feed in drought. Feed refusal, diarrhoea, liver abscessation.

Vitamin A deficiency - Slight eye problems – tears/ dust

Cold – 3 Growers not growing well during winter months and suffering from protein and energy (PED) due to poor feed intake (short days) die acutely from hypothermia/ glycaemia

## **Equines**

## **Mpumalanga**

### **Middelburg**

Tetanus - 1

## **Limpopo**

### **Modimolle**

Downers – 2 horses died, severe internal parasite infestation

## **Free State**

### **Parys**

Feed associated colic – 2

## **Eastern Cape**

### **Port Alfred**

Biliary – 1 (Kleinemonde)

Rain scald – 1 (Kleinemonde)

Sarcoid – 1 (Bathurst)

Diarrhoea – 1 Foal (Bathurst)

## **Western Cape**

### **Darling**

Retained after birth – 1

## **Northern Cape**

### **Colesberg**

Biliary - 2

## **Game**

## **Limpopo**

### **Bela-bela**

Lameness – Sable infection of the hoof, recovered

Lameness – Sable got caught up in the fence and injured achilles tendon

Trauma – buffalo – fighting between two bulls, one bull had wound caused by horn

### **Modimolle**

Sarcocystis – Impala ram dies one month after been moved

Actinobacillosis (Knopkakebeen) – Nyala ewe

## **North West**

### **Klerksdorp**

Red legged-ticks – 2 Zebra

## **KwaZulu-Natal**

## **Underberg**

Capture myopathy – 1 Eland

## **Eastern Cape**

### **Port Alfred**

Theileriosis – Sable 1

Veminosis – Sable 1

Diarrhoea – Sable calf 1

## **Swine**

## **Gauteng**

### **Onderstepoort**

Mange – 1

## **Dogs**

## **Northern Province**

### **Kathu**

Rabies - 1

## **Monthly report on Livestock and Wildlife isolations for August 2018 from Vetdiagnostix –Microbiology Laboratory, supplied by dr. Marijke Henton ([henton@vetdx.co.za](mailto:henton@vetdx.co.za))**

*Pasteurella multocida* [11] was the most common pathogen isolated from bovine respiratory tract infections, followed by *Mycoplasma* [7], *Mannheimia haemolytica* [4], *Histophilus somni* [4], *Trueperella pyogenes* [3] and *Mannheimia* biotype 8C [1] which is one of the biotypes associated with pneumonia.

Gangrenous myositis was caused by *Clostridium chauvoei* in 4 cases, and *C. novyi* in two.

Calf enteritis was associated in 13 cases with *E. coli*, and *Cryptosporidium* in 3. *Salmonella* Infantis was isolated on two different occasions from calves on the same farm. *Salmonella* Infantis is one of the pathogenic strains that cause disease from time to time in South Africa. *Salmonella* Infantis is considered to be an emerging pathogen.

*Salmonella* Dublin was isolated from a septicemic calf.

Enteritis in pigs was associated with *E. coli* in 5 cases.

Abscesses in cattle were associated with *Trueperella pyogenes* in 7 cases, in a goat in one case, and a kudu as well. Other causes of abscessation were a single case of *Corynebacterium pseudotuberculosis* in sheep, *Actinomyces bovis* in a goat, and *Actinomyces* sp in the joint of a lion.

A case of Blue Udder was due to *Staphylococcus aureus* in sheep. The Onderstepoort Biological Products Blue Udder vaccine contains two strains of *S. aureus*, and three of *Mannheimia [Pasteurella] haemolytica*, and should be effective in such cases.

**Feedlot report received from Drs. Shaun Morris and Eben du Preez for August 2018 ([edupreez1@telkomsa.net](mailto:edupreez1@telkomsa.net))**

Condition	Comments and Specie
Red gut	B 3
Blood gut	O 3
Botulism	B 2
Diarrhoea and deaths in lambs	O 3
Acidosis	B 3
Weak lambs, difficulty to adapt	O 3
Vitamin A deficiency with eye infection	O 3
Vitamin B 1 deficiency	B 2
Lungs	B 3

**Monthly report for Augustus 2018 from Dr R D Last (BVSc; M.Med.Vet(Path); MRCVS)**

**Specialist Veterinary Pathologist, Vetdiagnostix - Veterinary Pathology Services**

LIVESTOCK DISEASE SURVEILANCE			
LIVESTOCK SPECIES	DISEASE AGENT	NO. CASES	LOCATION
Bovine, Dairy calf	Cryptosporidiosis	1	Darling, W. Cape
Bovine, Dairy calf	Cryptosporidiosis	1	Dundee, KZN
Ovine, Lamb 3 weeks	Clostridial enterotoxaemia	1	Parys, Free State
Bovine, Dairy calf	Cryptosporidiosis	1	Darling, W. Cape
Bovine, Beef Calf	Cestrum poisoning	1	Howick, KZN

Bovine, Dairy Calf	<i>Pasteurella multocida</i> septicaemia	1	Dundee, KZN
Bovine, Dairy Calf	Cryptosporidiosis	1	Dundee, KZN

WILDLIFE DISEASE SURVEILLANCE			
WILDLIFE SPECIES	DISEASE AGENT	NO. CASES	LOCATION
White rhino, 3 month calf	Shuni virus encephalitis	1	Aliwal North, Northern Cape
White rhino, Adult cow	Iron storage disorder	1	Potchefstroom, North West
Buffalo, Adult Cow	Pulmonary hypertension	1	Rooiberg, Limpopo

**Monthly report on Livestock and Wildlife isolations for August 2018 from IDEXX Laboratories supplied by dr. Liza du Plessis ([Liza-DuPlessis@idexx.com](mailto:Liza-DuPlessis@idexx.com))**

Condition	Comments and Specie
Tapeworms	C 1
Coccidiosis	C 2
<i>E. coli</i>	B 1
Hepatotoxicity	B 1
Lungs	B,G 1
Abortion	B,O,C 1



Section of Pathology  
Department of Paraclinical Sciences  
Faculty of Veterinary Science



5<sup>th</sup> September 2018  
DAFF  
Import/Export Policy Unit Subdirector

**Monthly report: Faculty of Veterinary Science cases**  
**Cases sent to referring veterinarians between 27<sup>th</sup> July and 5<sup>th</sup> September 2018**

Cases from State vet Skukuza or Orpen

Cases imported with master permit (none)

PMDate	Species	Final	Histo No
17-Jul-18	African Buffalo	Acute renal failure	S2472-18
19-Jul-18	White Rhino	Cachexia	S2513-18
20-Jul-18	Burchell's Zebra	Suspected chronic peritonitis	S2545-18
25-Jul-18	Cheetah	Liver and renal failure, cholangitis	S2646-18
30-Jul-18	Lion	Suspected hypokalaemic polymyopathy	S2643-18
08-Aug-18	Impala	Emaciated, liver fibrosis	S2757-18
08-Aug-18	Roan	Bronchopneumonia	S2741-18
08-Aug-18	African Buffalo	Cardiac, skeletal myopathy	S2750-18

Kind regards,



Prof. Emily Mitchell